

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A23C 19/04, 19/032	A1	(11) International Publication Number: WO 00/54601 (43) International Publication Date: 21 September 2000 (21.09.00)
(21) International Application Number: PCT/DK00/00109 (22) International Filing Date: 14 March 2000 (14.03.00) (30) Priority Data: PA 1999 00368 16 March 1999 (16.03.99) DK (71) Applicant: NOVO NORDISK A/S [DK/DK]; Enzyme Business Patents, Novo Allé, DK-2880 Bagsværd (DK). (72) Inventor: NIELSEN, Per, Munk; Rytterstien 29A, DK-3400 Hillerød (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS FOR PRODUCING CHEESE (57) Abstract The present invention relates to a process for producing cheese from enzyme-treated cheese milk, and the use of the resulting produced cheese as ingredient in food products. More particularly, the present invention relates to a process for producing cheese from cheese milk treated with an enzyme selected from the group of phospholipases, in particular phospholipase A ₁ , A ₂ and B.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR PRODUCING CHEESE

TECHNICAL FIELD

5 The present invention relates to a process for producing cheese from enzyme-treated cheese milk, and the use of the resulting produced cheese as ingredient in food products. The present invention also relates to a process for stabilizing the fat in a milk composition.

10 BACKGROUND OF THE INVENTION

In cheese products, the state of the fat phase is important to the properties of the cheese. The fat phase is particularly important for the stabilisation of the cheese during production and ripening, but also for the final cheese to be used, eaten as
15 such, or used in prepared ready-to-eat dishes e.g. pizza, toast or burgers.

Also, the oiling-off properties of cheese products is an important quality parameter. Oiling-off is the tendency to form free oil upon storage and melting. Excessive oiling-off is a defect most often related to heated products wherein cheese is used, e.g.
20 pizza and related foods (cf. e.g. Kindstedt J.S; Rippe J.K. 1990, J Dairy Sci. 73: 867-873. It becomes more and more important to control/eliminate this defect, as the consumer concern about dietary fat levels increases. Free oil/fat in a product is perceived as a high fat content, and is generally undesirable.

25 In other food products the fat phase is often stabilised by mechanic emulsification, e.g. homogenisation. This technology is generally not applicable in cheese production as homogenisation of the cheese milk has a negative influence on the coagulation properties of the cheese milk and on the yield as well as the taste of the cheese produced therefrom.

30

In GB 1,525,929 it is disclosed as known to prepare stabilized oil-in-water emulsions using monoacyl glycerophosphatide obtained by subjecting diacyl

glycerophosphatide to the action of phospholipase A. GB 1,525,929 further describes use of phospholipase A treated phospholipoprotein-containing material for preparing oil-in-water emulsions, i.e. use of phospholipase treated material as an emulsion stabiliser for oil-in-water emulsions of which sauces, dressings and
5 mayonnaise is mentioned. Cheese is not disclosed in GB 1,525,929.

So-called lecithinase activity, disclosed as phospholipase activity, has been reported for bacterial contaminants in milk, as well as the use of such milk for cheese production: "J.J Owens, Observations on lecithinases from milk contaminants,
10 Process Biochemistry, vol. 13 no. 1, 1978, page 10-18" and "J.J Owens, Lecithinase Positive Bacteria in Milk, Process Biochemistry, vol. 13, page 13-15, 1978".

US 4,861,610 discloses a process for preparing a cheese composition, i.e. processed cheese, for incorporation into food material where monoacyl glycerophospholipid, fat, water and molten salt is added to cheese. The process comprises
15 a heating treatment to dissolve the cheese (the cheese being mixed with among others mono acyl glycerophospholipid) before the addition of fat. Subsequently the cheese-composition is emulsified by a mixer. US 4,861,610 does not disclose treatment of milk with phospholipase and manufacturing of cheese from the enzyme
20 treated milk.

There is a need for an improved process for the manufacturing of cheese, in particular a process for improving the stability of the fat in cheese.

25 SUMMARY OF THE INVENTION

The invention provides a process for producing cheese, which comprises the steps of:

- a) treating cheese milk or a fraction of cheese milk with a
30 phospholipase; and
- b) producing cheese from the cheese milk.

wherein step a) is conducted before and/or simultaneously with step b).

There is provided a process for improving the properties of cheeses; in particular the fat stability of cheese and cheese milk is improved by the present invention. The inventor has found that enzyme treatment of cheese milk significantly enhances the stability during a heat treatment of cheese produced from said phospholipase-treated cheese milk. By the process of the invention is also provided a method for increasing the yield in cheese production.

The invention further relates to the use of phospholipase in the manufacturing of cheese products, wherein the phospholipase treatment is conducted on the cheese milk or a fraction thereof before and/or during the production of the cheese. The invention also relates to cheeses obtainable, in particular obtained, by any of the processes described herein.

15 DETAILED DISCLOSURE OF THE INVENTION

The present invention relates to a process for producing cheese, which comprises the steps of:

- a) treating cheese milk or a fraction of cheese milk with a phospholipase; and
- 20 b) producing cheese from the enzyme-treated cheese milk of step a),
wherein step a) is conducted before step b) and/or simultaneously with step b).

Thus, step a) and b) of the process of the invention may be conducted simultaneously, i.e. the phospholipase reacts in the cheese milk at more or less the same time as the milk coagulant forms the coagulum.

Cheese milk and the production of cheese:

In the present context, the term "cheese" may be any kind of cheese and includes, e.g., natural cheese, cheese analogues and processed cheese. The cheese may be obtained by any suitable process known in the art, such as, e.g., by enzymatic coagulation of the cheese milk with rennet, or by acidic coagulation of the cheese milk with food grade acid or acid produced by lactic acid bacteria growth. In one

embodiment, the cheese manufactured by the process of the invention is rennet-curd cheese. Thus, in one embodiment, the cheese is manufactured with rennet. In step b) of the process of the invention the cheese milk may be subjected to a conventional cheese-making process. Rennet is commercially available, e.g. as Naturen® (animal rennet), Chy-max® (fermentation produced chymosin), Microlant® (Microbial coagulant produced by fermentation), all from Chr-Hansen A/S, Denmark).

Processed cheese may be manufactured from natural cheese or cheese analogues by cooking and emulsifying the cheese including emulsifying salts (e.g. phosphates and citrate), and may also include spices/condiments. In one embodiment, the cheese product of the process of the invention is not processed cheese.

By cheese analogues is understood cheese-like products, in which part of the composition is non-milk constituents, such as e.g. vegetable oil. Another example for cheese analogue is cheese base. The process of the present invention is applicable in producing cheese analogues as long as the product contains fat (e.g. milk fat, such as, e.g., cream) as a part of the composition.

The cheeses produced by the process of the present invention comprise all varieties of cheese, such as, e.g. Campesino, Chester, Danbo, Drabant, Herregård, Manchego, Provolone, Saint Paulin, Soft cheese, Svecia, Taleggio, White cheese, including rennet-curd cheese produced by rennet-coagulation of the cheese curd; ripened cheeses such as Cheddar, Colby, Edam, Muenster, Gryere, Emmenthal, Camembert, Parmesan and Romano; fresh cheeses such as Mozzarella and Feta; acid coagulated cheeses such as cream cheese, Neufchatel, Quarg, Cottage Cheese and Queso Blanco; and pasta filata cheese. One embodiment relates to the production of pizza cheese by the process of the invention.

In cheese manufacturing the coagulation of the casein in milk may be performed in two ways: the so-called rennet-curd and acid-curd cheese. In cheese production these two types of curds makes up two major groups of cheese types. Fresh acid-curd cheeses refer to those varieties of cheese produced by the coagulation of milk,

cream or whey via acidification or a combination of acid and heat, and which are ready for consumption once the manufacturing without ripening are completed. Fresh acid-curd cheeses generally differ from rennet-curd cheese varieties (e.g. Camberbert, Cheddar, Emmantal) where coagulation normally is induced by the action of rennet at pH values 6.4-6.6, in that coagulation normally occurs close to the isoelectric point of casein, i.e. e.g. at pH 4.6 or at higher values when elevated temperatures are used, e.g. in Ricotta pH 6.0 and 80°C. In a preferred embodiment of the invention, the cheese belongs to the class of rennet curd cheeses. In further embodiments the term cheese also includes acid curd cheese, including fresh acid-curd cheeses.

Mozzarella is a member of the so-called pasta filata, or stretched curd, cheeses which are normally distinguished by a unique plasticizing and kneading treatment of the fresh curd in hot water, which imparts the finished cheese its characteristic fibrous structure and melting and stretching properties, cf. e.g. "Mozzarella and Pizza cheese" by Paul S. Kindstedt, Cheese: Chemistry, physics and microbiology, Volume 2: Major Cheese groups, second edition, page 337-341, Chapman & Hall. Pizza cheese as used herein includes cheeses suitable for pizzas and they are usually pasta filata / stretched curd cheeses. In one embodiment, the process of the invention further comprises a heat/stretching treatment as for pasta filata cheeses, such as for the manufacturing of Mozzarella.

The cheese milk to be treated by the process of the present invention may comprise one or more of the following milk fractions: skim milk, cream, whole milk, buttermilk from production of sweet or acidified butter, whey protein concentrate, and butter or butter oil. The cheese milk to be phospholipase treated by the process of the invention may also comprise raw milk.

In further embodiments of the invention, the cheese milk to be phospholipase-treated is prepared totally or in part from dried milk fractions, such as, e.g., whole milk powder, skim milk powder, casein, caseinate, total milk protein or buttermilk powder, or any combination thereof.

The term "cheese milk", in particular in step b) of the process of the invention, is the milk-based composition from which the cheese is prepared. Thus, in the process of the invention the cheese may be produced from a milk-based composition ("the
5 cheese milk") of which all or only a portion has been subjected to a phospholipase treatment. The term "cheese milk" as used herein may encompass the term "fraction of the cheese milk" unless it is clear from the context that the two terms refer to different meanings.

10 The term "fraction of the cheese milk" in the context of the invention, in particular in step a) of the process of the invention, means the fraction of the cheese milk which is subjected to the enzymatic treatment of the invention. "The fraction of the cheese milk" may comprise one or more of the milk fractions as defined herein, i.e., e.g., skim milk, cream, whole milk, buttermilk from production of sweet or acidified butter,
15 whey protein concentrate, butter and butter oil. The butter may, e.g., be in a melted form. The fraction of the cheese milk to be treated may also comprise raw milk and it may also be prepared from dried milk fractions as already described herein. The term "fraction of the cheese milk" in the context of the present invention means one or more of the components of the cheese milk to be treated. When a fraction of the
20 cheese milk is phospholipase treated in step a) then step a) is performed before and not during step b). After the enzymatic treatment of the fraction of the cheese milk, the fraction is combined with one or more milk fractions to make of the cheese milk from which the cheese is prepared in step b).

25 The enzyme treatment in step a) may be conducted on a fraction of the cheese milk or it may be conducted on the cheese milk as such. Thus, within the scope of the invention is a process for producing cheese, which comprises the steps of: step i) treating a fraction of cheese milk with a phospholipase; step ii) preparing cheese milk from the treated fraction of step i); and step iii) producing cheese from the
30 cheese milk of step ii). It is also contemplated that in step ii) the enzyme treated cheese milk fraction of step i) may be combined with (a) non-phospholipase and/or (a) phospholipase treated cheese milk fraction(s) to provide the cheese milk from

which the cheese is produced in step iii). Step i) corresponds to step a); and step iii) corresponds to step b) as used herein.

In preferred embodiments, the cheese milk or the cheese milk fraction, which is to be
5 enzyme-treated, comprises or consists of cream. In further embodiments, the cheese
milk or the fraction of the cheese milk, which is to be enzyme-treated, comprises or
consists of butter. In still further embodiments, the cheese milk or the fraction of the
cheese milk, which is to be enzyme-treated, comprises or consists of buttermilk. In
one embodiment, the enzyme treated milk of step a) is not dried before step b). In
10 further embodiments, the process of the invention does not include a particular step
for lowering the total fat content of the cheese, such as, e.g., the process disclosed
in EP 531 104 A2 which relates to a process for reducing the lipid content in food.

Milk from different species of animals may be used in the production of cheese.
15 Thus, "milk" may be the lacteal secretion obtained by milking, e.g., cows, sheep,
goats, buffaloes or camels.

The milk for production of cheese may be standardised to the desired composition
by removal of a portion or of all of any of the raw milk components and/or by adding
20 thereto additional amounts of such components. This may be done by separation of
the raw milk into cream and skim milk at arrival to the dairy. Thus, the cheese milk
may be prepared as done conventionally by fractioning the raw milk and recombining
the fractions so as to obtain the desired final composition of the cheese milk. The
separation may be made in continuous centrifuges leading to a skim milk fraction
25 with very low fat content (i.e. e.g. < 0.5%) and cream with e.g. > 35% fat. The
"cheese milk" may be composed by mixing cream and skim milk. In a preferred
embodiment the cheese milk or the fraction of the cheese milk to be treated with
phospholipase is not derived from cheese.

30 The cheese milk, including the cheese milk fraction, to be treated with
phospholipase comprises phospholipids, such as e.g. lecithin. The cheese milk may
have any total fat content which is found suitable for the cheese to be produced by

the process of the invention, such as, e.g., about 25% fat (of dry matter), such as e.g. in the range 10-50% fat, of which, e.g., about 0.06% is phospholipids, such as e.g. 0.02-5% (w/w) of the total fat content is phospholipids.

5 Conventional steps may be taken to secure low bacterial counts in the cheese milk. It is generally preferred not to pasteurise the skim milk because heat denatured proteins in the cheese milk has a negative influence on the coagulation of the milk, and retards the ripening of the cheese. The bacterial count of the skim milk fraction may thus be lowered by other technologies, as for instance microfiltration or
10 bactofugation. The cream may be pasteurised to get the low bacterial count in the product.

The process of the invention may further comprise the step of c) subjecting the treated cheese milk or the cheese milk fraction to a heating treatment after step a)
15 and before step b). In one embodiment, the process of the invention comprises the steps of: step i) treating a fraction of cheese milk (such as, e.g., cream) with a phospholipase; step ii) subjecting the enzyme treated fraction of step i) to a heating treatment; step iii) combining the milk fraction of step ii) with (a) non-enzyme and/or (a) phospholipase treated milk fraction(s) to obtain the cheese milk from which the
20 cheese is produced; and step iv) producing cheese from the cheese milk of step iii), wherein step i) corresponds to step a); and step iv) corresponds to step b) as used herein. Accordingly, in one embodiment the process of the invention, wherein the fraction of cheese milk is cream, may further comprise the step of subjecting the cream to pasteurization after step a) and before step b).

25

In the process of the invention the cheese milk or the fraction of the cheese milk may be subjected to a homogenization process before the production of cheese, such as e.g. in the production of Danish Blue Cheese. The homogenization may be applied before and/or after the treatment with the phospholipase. In other embodiments,
30 however, the cheese milk in step b) is not subjected to a homogenization process before the production of cheese.

The enzymatic treatment:

The enzymatic treatment in the process of the invention may be conducted by dispersing the phospholipase into cheese milk or a fraction of the cheese milk, and allowing the enzyme reaction to take place at an appropriate holding-time at an
5 appropriate temperature. The treatment with phospholipase may be carried out at conditions chosen to suit the selected enzymes according to principles well known in the art. The enzymatic treatment is a treatment in which the milk fat fraction of the cheese milk is treated with phospholipase.

- 10 The enzymatic treatment may be conducted at any suitable pH, such as e.g. in the range 2-10, such as at a pH of 4-9 or 5-7. It may be preferred to use pH of 5.5-7.0.

The process of the invention may be conducted as a phospholipase treatment of cheese milk or a fraction of the cheese milk during cold storage at 3-7°C, e.g. for at
15 least 2 hours, e.g. in the range of 2-48 hours, or at least 5 hours, e.g. 5-24 hours.

The process may also be conducted so that the phospholipase is allowed to react at coagulation conditions 30-45°C (e.g. for at least 5 minutes, such as, e.g., for at least 10 minutes or at least 30 minutes, e.g. for 5-60 minutes) during, e.g., the cheese making process of step b). Further, the process may be conducted so that before
20 coagulation of the cheese milk, the phospholipase is allowed to react on a milk fraction, e.g. cream, at the temperature optimum for the phospholipase, e.g. at 45-80°C, such as 47-80°C, or 50-80°C, e.g. for at least 10 minutes, such as at least 30 minutes, e.g. in the range of 10-180 minutes.

- 25 Optionally, after the enzymatic treatment the phospholipase enzyme protein is removed/reduced and/or the enzyme is inactivated.

A suitable enzyme dosage will usually be in the range of 0,01-1% (w/w) of the fat content, such as, e.g., 0.1 - 1.0 %, particularly 0.2 % (w/w) corresponding to 2000 IU
30 per 100 g fat. One IU (International Unit) is defined as the amount of enzyme producing 1 micro mole of free fatty acid per minute under standard conditions: Egg yolk substrate (approximately 0.4% phospholipids), pH 8, 40°C, 6mM Ca⁺⁺,

Analytical method AF 280 available on request from Novo Nordisk A/S and is described in the Examples. The enzyme dosage is based on w/w fat content of the treated cheese milk such as cream illustrated in the examples. Alternatively, the enzyme dosage may be determined by the other assays as described herein.

5

The enzymatic treatment may be conducted batchwise, e.g. in a tank with stirring, or it may be continuous, e.g. a series of stirred tank reactors.

In one embodiment, the phospholipase is added to the cream fraction to carry out a
10 separate phospholipase treatment of this fraction at a temperature in the range 45-80°C. In further embodiments, the phospholipase is added immediately before, or at the same time as the cheese rennet, e.g., at 32-36°C.

Enzymes to be used in the process of the invention:

15 The enzyme used in the process of the present invention include a phospholipase, such as, phospholipase A₁, phospholipase A₂ and phospholipase B. In the process of the invention the phospholipase treatment may be provided by one or more phospholipase, such as two or more phospholipases, e.g. two phospholipases, including, without limitation, treatment with both type A and B; both type A₁ and A₂;
20 both type A₁ and B; both type A₂ and B; or treatment with two different phospholipase of the same type. Included is also treatment with one type of phospholipase, such as A₁, A₂ or B.

Phospholipids, such as lecithin or phosphatidylcholine, consist of glycerol esterified
25 with two fatty acids in an outer (sn-1) and the middle (sn-2) positions and esterified with phosphoric acid in the third position; the phosphoric acid, in turn, may be esterified to an amino-alcohol. Phospholipases are enzymes which participate in the hydrolysis of phospholipids. Several types of phospholipase activity can be distinguished, including phospholipases A₁ and A₂ which hydrolyze one fatty acyl
30 group (in the sn-1 and sn-2 position, respectively) to form lysophospholipid; and lysophospholipase (or phospholipase B) which can hydrolyze the remaining fatty

acyl group in lysophospholipid. Thus, the invention relates to use of enzymes that has the ability to hydrolyze one and/or both fatty acyl groups in a phospholipid.

Phospholipase A₁ is defined according to standard enzyme EC-classification as EC

5 3.1.1.32.

Official Name: Phospholipase A₁.

Reaction catalyzed:

phosphatidylcholine + H₂O <>

2-acylglycerophosphocholine + a fatty acid anion

10 Comment(s)

has a much broader specificity than ec 3.1.1.4.

Phospholipase A₂ is defined according to standard enzyme EC-classification as EC

3.1.1.4

Official Name: phospholipase A₂.

15 Alternative Name(s): phosphatidylcholine 2-acylhydrolase.

lecithinase a; phosphatidase; or phosphatidolipase.

Reaction catalysed:

phosphatidylcholine + h₂o <>

1-acylglycerophosphocholine + a fatty acid anion

20 comment(s): also acts on phosphatidylethanolamine, choline plasmalogen and phosphatides, removing the fatty acid attached to the 2-position.

The term "Phospholipase A" used herein in connection with an enzyme of the invention is intended to cover an enzyme with Phospholipase A₁ and/or
25 Phospholipase A₂ activity.

"Phospholipase B": Phospholipase B is defined according to standard enzyme EC-classification as EC 3.1.1.5.

Official Name: lysophospholipase.

30 Alternative Name(s): lecithinase b; lysolecithinase;

phospholipase b; or plb.

Reaction catalysed:

2-lysophosphatidylcholine + h(2)o <> glycerophosphocholine + a fatty acid anion

The term "phospholipase" used herein in connection with an enzyme of the invention is intended to cover enzymes which has enzyme activity towards phospholipids as defined herein. The term phospholipase as used herein, includes enzymes with phospholipase activity, i.e., e.g. phospholipase A (A₁ or A₂) or phospholipase B activity. The phospholipase activity may be provided by enzymes having other activities as well, such as e.g. a lipase with phospholipase activity. The phospholipase activity may e.g. be from a lipase with phospholipase side activity. In other embodiments of the invention the phospholipase enzyme activity is provided by an enzyme having essentially only phospholipase activity and wherein the phospholipase enzyme activity is not a side activity. In one embodiment of the invention, the phospholipase is not lipases having phospholipase side activity as defined in WO 98/26057.

The phospholipase may be of any origin, e.g. of animal origin (such as, e.g. mammalian), e.g. from pancreas (e.g. bovine or porcine pancreas), or snake venom or bee venom. Alternatively, the phospholipase may be of microbial origin, e.g. from filamentous fungi, yeast or bacteria, such as the genus or species *Aspergillus*, e.g. *A. niger*, *Dictyostelium*, e.g. *D. discoideum*; *Mucor*, e.g. *M. javanicus*, *M. mucedo*, *M. subtilissimus*; *Neurospora*, e.g. *N. crassa*; *Rhizomucor*, e.g. *R. pusillus*; *Rhizopus*, e.g. *R. arrhizus*, *R. japonicus*, *R. stolonifer*; *Sclerotinia*, e.g. *S. libertiana*; *Trichophyton*, e.g. *T. rubrum*; *Whetzelinia*, e.g. *W. sclerotiorum*; *Bacillus*, e.g. *B. megaterium*, *B. subtilis*; *Citrobacter*, e.g. *C. freundii*; *Enterobacter*, e.g. *E. aerogenes*, *E. cloacae* *Edwardsiella*, *E. tarda*; *Erwinia*, e.g. *E. herbicola*; *Escherichia*, e.g. *E. coli*; *Klebsiella*, e.g. *K. pneumoniae*; *Proteus*, e.g. *P. vulgaris*; *Providencia*, e.g. *P. stuartii*; *Salmonella*, e.g. *S. typhimurium*; *Serratia*, e.g. *S. liquefaciens*, *S. marcescens*; *Shigella*, e.g. *S. flexneri*; *Streptomyces*, e.g. *S. violaceoruber*; *Yersinia*, e.g. *Y. enterocolitica*. Thus, the phospholipase may be fungal, e.g. from the class *Pyrenomycetes*, such as the genus *Fusarium*, such as a strain of *F. culmorum*, *F. heterosporum*, *F. solani*, or a strain of *F. oxysporum*. The phospholipase may also

be from a filamentous fungus strain within the genus *Aspergillus*, such as a strain of *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus niger* or *Aspergillus oryzae*. A preferred phospholipase is derived from strain of *Fusarium*, particularly *F. oxysporum*, e.g. from strain DSM 2627 as described in WO 98/26057, especially described in claim 36 and SEQ ID NO. 2 of WO 98/26057. In further
5 embodiments, the phospholipase is a phospholipase as disclosed in PCT/DK/00664 (Novo Nordisk A/S, Denmark).

The phospholipase used in the process of the invention may be derived or
10 obtainable from any of the sources mentioned herein. The term "derived" means in this context that the enzyme may have been isolated from an organism where it is present natively, i.e. the identity of the amino acid sequence of the enzyme are identical to a native enzyme. The term "derived" also means that the enzymes may have been produced recombinantly in a host organism, the recombinant produced
15 enzyme having either an identity identical to a native enzyme or having it a modified amino acid sequence, e.g. having one or more amino acids which are deleted, inserted and/or substituted, i.e. a recombinantly produced enzyme which is a mutant and/or a fragment of a native amino acid sequence. Within the meaning of a native enzyme are included natural variants. Furthermore, the term "derived" includes
20 enzymes produced synthetically by e.g. peptide synthesis. The term "derived" also encompasses enzymes which have been modified e.g. by glycosylation, phosphorylation etc., whether in vivo or in vitro. The term "obtainable" in this context means that the enzyme has an amino acid sequence identical to a native enzyme. The term encompasses an enzyme that has been isolated from an organism where it
25 is present natively, or one in which it has been expressed recombinantly in the same type of organism or another, or enzymes produced synthetically by e.g. peptide synthesis. With respect to recombinantly produced enzyme the terms "obtainable" and "derived" refers to the identity of the enzyme and not the identity of the host organism in which it is produced recombinantly.

30

Accordingly, the phospholipase may be obtained from a microorganism by use of any suitable technique. For instance, a phospholipase enzyme preparation may be

obtained by fermentation of a suitable microorganism and subsequent isolation of a phospholipase preparation from the resulting fermented broth or microorganism by methods known in the art. The phospholipase may also be obtained by use of recombinant DNA techniques. Such method normally comprises cultivation of a host
5 cell transformed with a recombinant DNA vector comprising a DNA sequence encoding the phospholipase in question and the DNA sequence being operationally linked with an appropriate expression signal such that it is capable of expressing the phospholipase in a culture medium under conditions permitting the expression of the enzyme and recovering the enzyme from the culture. The DNA sequence may also
10 be incorporated into the genome of the host cell. The DNA sequence may be of genomic, cDNA or synthetic origin or any combinations of these, and may be isolated or synthesized in accordance with methods known in the art.

Suitable phospholipases are available commercially. As typical examples of the
15 enzymes for practical use, pancreas-derived phospholipase A2 such as Lecitase® (manufactured by Novo Nordisk A/S) is preferably used.

In further embodiments, the source of the phospholipase in the process of the invention, is from expressing the enzyme by the starter organism used in the
20 production of the cheese, such as e.g. by over-expressing the phospholipase in a lactic acid bacterium, including e.g. *Lactobacillus*. Alternatively, the treatment of the cheese milk or the fraction of cheese milk is conducted by the addition of the phospholipase, optionally in combination with the phospholipase provided from the starter culture as described herein.

25

In a preferred embodiment, the phospholipase is not obtained from microbial milk contaminants. Accordingly, in the process of the invention, the phospholipase enzyme treatment is not provided by the enzymatic action of phospholipase expressed by a microbial milk contaminant present in the milk or the milk fraction, at
30 least not a major portion. The microbial milk contaminants may be one or more of the group consisting of *Bacillus cereus*, *Bacillus cereus* var. *mycoides*, *Pseudomonas* sp., *Enterobacter liquifaciens* (*Klebsiella cloacae*), *Alcaligenes viscolactis*, *Corynebacterium*

rod. In one embodiment, the phospholipase is not obtained from these contaminants. In further embodiments the phospholipase is not identical to the enzymes disclosed in "J.J Owens, Observations on lecithinases from milk contaminants, Process Biochemistry, vol. 13 no. 1, 1978, page 10-18" and "J.J Owens, Lecithinase Positive
5 Bacteria in Milk, Process Biochemistry, vol. 13, page 13-15, 1978".

In the process of the invention, the phospholipase treatment may be performed by contacting the cheese milk and/or the cheese milk fraction with a purified phospholipase. The term "purified" as used herein covers phospholipase enzyme
10 protein free from components from the organism from which it is derived. The term "purified" also covers phospholipase enzyme protein free from components from the native organism from which it is obtained, this is also termed "essentially pure" phospholipase and may be particularly relevant for phospholipases which are naturally occurring phospholipases and which have not been modified genetically,
15 such as by deletion, substitution or insertion of one or more amino acid residues.

Accordingly, the phospholipase may be purified, viz. only minor amounts of other proteins being present. The expression "other proteins" relate in particular to other enzymes. The term "purified" as used herein also refers to removal of other
20 components, particularly other proteins and most particularly other enzymes present in the cell of origin of the phospholipase. The phospholipase may be "substantially pure", i.e. free from other components from the organism in which it is produced, i.e., e.g., a host organism for recombinantly produced phospholipase. Preferably, the enzymes are at least 75% (w/w) pure, more preferably at least 80%, 85%, 90% or
25 even at least 95% pure. In a still more preferred embodiment the phospholipase is an at least 98% pure enzyme protein preparation. In other embodiments the phospholipase is a phospholipase not naturally present in milk.

The term phospholipase includes whatever auxiliary compounds that may be
30 necessary for the enzyme's catalytic activity, such as, e.g. an appropriate acceptor or cofactor, which may or may not be naturally present in the reaction system.

The phospholipase may be in any form suited for the use in question, such as e.g. in the form of a dry powder or granulate, a non-dusting granulate, a liquid, a stabilized liquid, or a protected enzyme. Granulates may be produced, e.g. as disclosed in US 4,106,991 and US 4,661,452, and may optionally be coated by methods known in the art. Liquid enzyme preparations may, for instance, be stabilized by adding stabilizers such as a sugar, a sugar alcohol or another polyol, lactic acid or another organic acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

10 By the process of the invention, the lecithin content of the cheese may be reduced by at least 5%, such as at least 10%, at least 20%, at least 30%, at least 50%, such as in the range of 5-95% compared to a similar cheese making process but without the enzymatic treatment in step a).

15 In cow milk the lecithin constitutes normally more than 95% of the phospholipids in milk whereas the lysolecithin is approximately 1% of the phospholipids. Although the phospholipids represent normally less than 1% of the total lipids in cow milk, they play a particularly important role, being present mainly in the milk fat globule membrane. By the process of the present invention the lecithin content in the cheese
20 of step b) may be less than 90%, such as e.g. less than 80%, e.g. less than 60% or less than 50% of the total content of phospholipid in the cheese. In other embodiments of the invention the lysolecithin content in the cheese constitutes at least 5%, such as at least 10%, at least 20%, at least 30%, at least 50%, such as, e.g. in the range 5-99%, e.g. 5-90%, 10-90%, or 30-90% or 40-80% of the total
25 content of phospholipids in the cheese. The lecithin or lysolecithin content may be measured by any method known by the skilled person, e.g. by HPLC.

In a preferred embodiment, it is to be understood that the relative amount of lecithin to lysolecithin in the cheese produced is provided by conversion of lecithin to
30 lysolecithin by the treatment of the cheese milk or cheese milk fraction with phospholipase. In cheese the fat content is generally 65% (w/w), such as in the range of 10-60%.

The invention also relates to the use of phospholipase in the process of the invention. Accordingly, one embodiment relates to the use of a phospholipase in the manufacturing of a cheese product, comprising adding the phospholipase to cheese
5 milk or a fraction of the cheese milk, and processing the cheese milk to make the cheese. Thus, within the scope of the invention is the use of a phospholipase in the manufacturing of a cheese product, wherein the phospholipase treatment is conducted on cheese milk or a fraction thereof before and/or during the production of the cheese.

10

The present invention further relates to use of the cheese produced by the process of the invention in pizza, ready-to-eat dishes, processed cheese or as an ingredient in other food products. Accordingly, the cheese produced according to the process of the invention may be used in further processed food products like processed
15 cheese, pizza, burgers, toast, sauces, dressings, cheese powder or cheese flavours.

In further embodiments, the process of the invention further comprises the step of subjecting the cheese of step b) to a heating treatment, such as, e.g. in the range 150-350°C.

20

The invention also relates to a cheese obtainable, in particular obtained, by the process of the invention.

In a preferred embodiment, the invention relates to a process for improving the
25 stability of the fat in cheese, the process being as described herein. In one embodiment, the cheese manufactured by the process of the invention has a lowered diffusion of fat/oil, such as, e.g., a decrease in the "oily" diameter of at least 5%, such as at least 10%, at least 20%, at least 40%, e.g. a decrease of the "oily" diameter in the range 20-800%, e.g. 20-600%; the "oily diameter" being measured
30 as defined herein in Example 1 or 2.

In one embodiment, the invention relates to a process of the invention for increasing the yield of cheese, such as an increase in cheese fat yield and/or cheese protein yield in cheese production. Accordingly, by the present invention is provided a cheese manufacturing process leading to higher yield of the cheese and better stability of the fat phase of the cheese produced by the process of the invention. Thus, one embodiment relates to a process of the invention for improving the fat stability of cheese and/or increasing the yield in cheese production. The increase in fat-yield by the process of the invention may be at least 0,5%, such as e.g. in the range 0,5-10%, such as in the range of 0,5%-5%, measured e.g. as "fat by difference" as described in Example 3.

Further aspects of the invention:

The invention also provides a process for stabilizing the fat of a milk composition, which comprises treating a milk or a fraction of the milk with an enzyme selected from the group of phospholipases. The inventor has found that the emulsion stability of a milk composition can be improved by treating it with phospholipase. Thus, the invention also relates to a process for stabilizing the fat emulsion of a milk composition, which comprises treating a milk or a fraction of the milk with a phospholipase. The process may be conducted as in step a) of the process described herein.

By the present invention is also provided a method for producing improved UHT cream, in particular having improved fat stability, which process for producing UHT-cream comprises the steps of: step a) treating cream with a phospholipase; and step b) subjecting the phospholipase treated cream of step a) to UHT treatment, wherein step a) is performed before step b). The invention further relates to UHT-cream obtainable or obtained from such process of the invention. Finally, the invention relates to the use of a phospholipase in the manufacturing of UHT-cream.

By the present invention is also provided a process for producing a cream liquor, which comprises the steps of: step a) treating a milk composition (such as, e.g., cream) with a phospholipase; and step b) producing a cream liquor from the enzyme

treated milk composition. The invention further relates to cream liquor obtainable or obtained from such process of the invention. Finally, the invention relates to the use of a phospholipase in the manufacturing of cream liquor.

- 5 The present invention is further illustrated in the following examples which is not to be in any way limiting to the scope of protection.

EXAMPLES

10 Determination of phospholipase activity:

The determination of the phospholipase activity may be made according to the principles described in the following. In this method is used egg yolk as a lecithin-rich substrate.

- Principle: pH stat titration. Homogenized egg yolk containing phospholipids is
15 hydrolyzed by phospholipase in the presence of calcium and sodium deoxycholate at pH 8.0 and 40°C in a pH-stat. The released fatty acids are titrated with 0.1 N sodium hydroxide and base volume is monitored as a function of the time. One phospholipase unit is defined as the enzyme quantity which under standard conditions produces one micro-equivalent free fatty acid.

20 Reagents:

0.016 M Sodium deoxycholate ($C_{24}H_{39}NaO_4$), 0.32 M Calcium chloride,
0.01 N Hydrochloric acid, 0.1 N Sodium hydroxide.

- Substrate: Add one egg yolk to 100 ml water and homogenize in a disperser. Filter the homogeneous substrate through double gauze. Add 5 ml of 0.32 M calcium
25 chloride to stabilize the filtrate.

Titrate mixture: Mix 100 ml of substrate with 50 ml of 0.016 sodium deoxycholate.

- Alternatively, the following assays may be used for qualitative or quantitative determination of phospholipase activity as described in DK 99/00664 (Novo Nordisk
30 A/S, Denmark): "Phospholipase activity (PHLU)", "Phospholipase activity (LEU)", "Phospholipase monolayer assay", "Plate assay 1" and "Plate assay 2" or as described in WO 98/26057 (Novo Nordisk A/S): the "NEFA-C test".

EXAMPLE 1 PHOSPHOLIPASE TREATMENT OF CREAM FOR CHEESE PRODUCTION

- 5 In this example cream was treated with phospholipase before combined with skim milk to prepare the cheese milk.

Raw materials

Composition of the cheese milk:

10 Skim milk	0.1 % fat	1820 ml
cream	38.0 % fat	180 ml
Butter milk (comprising the starter culture)		50 ml
CaCl ₂		0.4 g

Enzymes:

- 15 1) Rennet (Acid aspartic *Rhizomucor miehei* protease – EC 3.4.23.6) dosage: 0.107g
2) Lecitase® (pancreas-derived phospholipase A₂ obtainable Novo Nordisk A/S),
dosage(based on fat):0,2 % (w/w)

Cheese production

- 20 Method: The cream was treated separately with phospholipase (Lecitase®, manufactured by Novo Nordisk A/S, Denmark; dosage 0.2 % (based on w/w fat content)) by incubating the mixture in a 50°C water bath for 30 minutes. The treated cream was mixed with the skim milk to a total fat content of 3.5 % (w/w) of the mix, and placed in a 33-35°C water bath. CaCl₂ was added to the 2 liter cheese milk, and
25 starter culture (i.e. butter milk) was added, and the mixture was stirred for 5-10 minutes. The milk was left for 30 minutes without any stirring. Then, rennet (acid aspartic *Rhizomucor miehei* protease) was added, and the milk was stirred for 1 minute. Subsequently, the clotting point was defined (approximately 12 minutes), and the milk was standing for about 25-26 minutes before cutting.

30

The clotting point (clotting time) is the time passed from the addition of rennet to the first sign of flocculation, and is determined by moving a black rod in the milk i.e. the

clotting point is the point when the first visible precipitation of paracasein is observed on the rod.

The actual cutting time was defined by doing a test as follows: With a clotting stick a
5 small cut was made on the surface of the cheese, the stick was putted under the cut
and moved forward, when the cheese was separated with two sharp edges, and the
whey would collect in between the two edges then the cheese was ready for cutting.

The cheese was stirred gently with a whisk to break it up, and was left for two
10 minutes. After the two minutes, the cheese was occasionally stirred during 10-15
minutes to separate the whey from the curd. The steadily appearing whey and the
curd were transferred to a sieve containing a cloth. The whey drained away through
the cloth for 1-2 hours. A 0.6 kg weight was placed on the top of the cheese to
remove more whey, and thereafter the cheese was stored overnight at room
15 temperature.

Result: The cheese production data (clotting time, cutting time, amount of whey,
protein in whey and amount of cheese) are presented in table 1.

20 **Melting of cheese**

Method: Before melting, the cheese was cut into pieces with a height of about 2.5 cm
and diameter of about 8 cm. The cheese samples was heated in a oven at 250°C for
8 minutes. The diameter of the cheese after heating was measured as the average
of two diameters.

25 Result: cf. Table 1 below.

Diffusion of fat/oil

Method: Before the diffusion test was made, the cheese was cut into pieces with a
height of about 2.5 cm and diameter of about 8 cm. The diffusion of fat/oil was
30 measured after heating the cheese samples in an oven at 200°C for 3 minutes. The
diameter on the liquid diffused out on a filter paper - Whatman 40 - after heating was
measured as the average of four diameters.

Result: cf. Table 1 below.

The measurements of samples obtained in Example 1 are shown in table 1

The trials were conducted as two set of trials, where different batches of milk/cream
5 raw materials were used. One set is the column 1 and 2. The second set is column
3,4 and 5.

TABLE 1. Trials with phospholipase used for treatment of cream in cheese production.

Trial no.	1* Control	2 with Lecitase	3* Control	4 with Lecitase	5 with Lecitase
Data from Cheese production					
Clotting, minutes	13 ³⁰	14 ¹⁵	16 ²⁰	17 ⁰⁰	16 ⁰⁰
Cutting, minutes	31 ³⁰	31 ³⁰	35 ³⁰	35 ³⁰	34 ³⁰
Weight of whey	1635.5	1660.1	1625.8	1645.8	1668.7
Weight of cheese	314.75	296.74	327.44	322.75	318.77
% protein in whey	0.96	0.92	0.84	0.83	0.83
g protein in whey	15.7	15.3	13.7	13.7	13.9
Result melting of cheese					
Average diameter after heating, cm	8.0	7.5	8.0	7.3	7.3
Result diffusion of fat/oil					
"Oily" diameter on filter paper after heating, cm	6.6	5.8	6.8	5.4	5.4

10 * Trials 1 and 3, no Lecitase[®] added. Trial 1* is the control for trial no. 2 and Trials 3* is the control for trial no. 4-5.

From the Result Table (cf. "melting of cheese" and "diffusion of fat/oil" shown at the bottom of Table 1) it clearly appears that cheese made with an initial phospholipase
15 treatment of the cream improves the stability of the cheese during a heat treatment. This is seen from the measurements of the average diameter of sample after heating, which is smaller when the cream for the cheese milk has been treated with phospholipase. Further, the oily diameter on the filter paper is reduced when treating the cream with phospholipase.

The results demonstrate that it is possible to obtain a stability improvement of cheese during heat treatment with the process of the invention.

5 **EXAMPLE 2 PHOSPHOLIPASE TREATMENT OF CREAM TO TEST THE STABILISING EFFECT ON THE CREAM**

Method: Cream (200 g) was incubated 15 minutes at 50°C with 3 different dosages of Lecitase® 10L, (Novo Nordisk A/S, Denmark) (dosage based on estimated fat =
10 36%):

1. Dosage 0 %
2. Dosage 0.2 % (= 0.0725% of the cream)
3. Dosage 1.0 % (= 0.36% of the cream)

After the incubation the cream was cooled to 5°C. The cream was whipped in a
15 standardised way (Philips mixer 5 speed, the first 2 minutes run at speed 4, and speed 5 the rest of the time). 100 g whipped cream was filled into a funnel and allow to stand dripping for 1 hour at room temperature.

Samples and results:

20 Sample 1 was whipped for 7 minutes and 25 seconds. About 300 ml cream. Amount of dripping after 1 hour in funnel = 2,5 ml.

Sample 2 was whipped 20 minutes and no foam formation was observed. The sample became a little bit thicker, and obtained a light yellow colour.

25

Sample 3 was whipped for 15 minutes and no foam formation was observed. The sample became a little bit thicker and obtained a light yellow colour.

The ability to foam was destroyed by Lecitase® treatment. The foam formation in
30 cream is based on collapse of fat globules, and formation of a continuous fat phase forming the foam. In accordance with the process of the present invention the above

results show that Lecitase[®] treatment improves emulsification and stabilisation of the fat in the cream.

EXAMPLE 3 PHOSPHOLIPASE TREATMENT OF CREAM FOR 5 MOZZARELLA CHEESE PRODUCTION.

In this example cream was treated with phospholipase before being combined with skim milk to produce the cheese milk.

10 Raw materials

Composition of the cheese milk:

Skim milk, pasteurised	0.83 % fat	13.63L
cream	30 % fat	1.37 L
CaCl ₂		0.4 g/2 kg cream

- 15 Starter cultures LH100 and TA061 (Lactic acid bacteria) from Rhodia Foods (Rhodia Inc, Madison, WI, USA) 0.6g of each.

Enzymes:

- 1) Rennet (Acid aspartic *Rhizomucor miehei* protease – EC 3.4.23.6) dosage w 50
20 KRU/g (KRU method obtainable from Novo Nordisk A/S: 0.60 g
2) Lecitase[®] (pancreas-derived phospholipase A₂ obtainable Novo Nordisk A/S),
dosage(based on fat):0,2 % (w/w)

Cheese production

- 25 Method: The cream was treated separately with phospholipase (Lecitase[®],
manufactured by Novo Nordisk A/S, Denmark; dosage 0.2 % (based on w/w fat
content)) by incubating the mixture in a 50°C water bath for 30 minutes with CaCl₂.
The treated cream was mixed with the skim milk to a total fat content of 3.5 %, and
placed in 15 L cheese vat (using a cheese unit with 2x15L vats obtainable from GEA
30 Liquid processing, Haderslevvej 36, 6000 Kolding, Denmark). The milk was
equilibrated to 34.4 °C, and starter culture was added, and the mixture was gently
agitated for 4-5 minutes before the rennet was added. Then, rennet (acid aspartic

Rhizomucor miehei protease) was added, and the milk was stirred for 3 minutes. Subsequently the milk was standing for about 35 minutes before cutting

The actual cutting time was defined by doing a test as follows: With a clotting stick a
5 small cut was made on the surface of the cheese - then the stick was putted under the cut and moved forwards so as the cheese was separated with two sharp edges, and the whey would collect in between the two edges. The cutting was performed with ½ inch knives.

10 The cutted curd was then heated to 41.1 °C (takes about 30 min). The cheese was then agitated gently until the curd pH reached pH 5.90, whereafter the whey was drained and piled to a 5 cm mat. The curd is cut into 1.5 inch cubes when the pH reach 5.25 and covered in cold tap water for 15 min. The curd is weighed after tap water is drained of. NaCl (0.2 % of the cheese milk weight) was added dry to the
15 cubed curd in 3 portions. The curd was then stretched at 63 °C at 9 rmp by twin screw stretcher (Supreme Micro Mixing Machine obtained from Stainless Fabricating Inc., Columbus. WI, USA), where after the stretched curd is placed in 7 °C water for 30 min and 7 °C F brine (23% NaCl) for another 90 minutes.

20 The cheese production data (Protein in cheese, measured by Dumas Combustion Method (LECO) moisture measured by CEM Automatic Volatility Computer, model AVC-80 from CEM Corp., Matthews, NC, 28108, and fat by difference) are collected in table 2.

25 Result: cf. Table 2 below.

Diffusion of fat/oil

Method: The diffusion of fat/oil was tested on cheese samples heated in the oven at 90°C for 5 minutes. Before the diffusion test was made, the cheese was ground in an
30 osterizer blender to achieve uniformity of sample. Subsequently 2.0 grams were molded into a metal ring (2 cm) and placed in the center of a Whatman # 4 filter .

The oiling off was determined by difference in the areas between the ring of oil and the circle of the cheese (triplicate determinations). Result: cf. Table 2 below.

Meltability

- 5 Method: squeeze-flow rheometry, ref. Ak, M.M., and Gunasekaran. 1995. J. Texture Stud. 26: 695-711.

Parameters: 0.1 mm/ sec crosshead speed on Stable Micro Systems, TA-XT2 texture analyzer. Cylindrical samples diameter 25 mm, height 15 mm) were placed in 50°C water bath for 5 min. to equilibrate temperature.

10

The meltabilities of the P-lipase treated samples were only slightly lower than their respective controls.

Measurements of samples obtained in examples 3 are shown in table 2

15

The trials were conducted as two set of trials, where different batches of milk/cream raw materials were used.

20 TABLE 2. Data for Example 3: Trials with phospholipase used for treatment of cream in cheese production.

Trial no.	1	1-control	2	2 control
Data from cheese production				
Protein in cheese	18.8%	18.1%	20.4%	20.5%
Moisture	43.1%	45.1%	46.3%	46.9%
Fat by difference	38.1%	36.8%	33.3%	33.6%
Result from oiling off "Oily" area on filter paper after heating, in percent of the cheese area	57%	176%	<10%	61%

* Trials 1 and 2 is with Lecitase® added. Trial 1-control and 2-control are the corresponding trials run in parallel but without Lecitase®. In experiment 1 the oiling of was measured after 5 days in refrigeration, and in experiment 2 the oiling of was measured after storage in refrigeration for 8 days .

From the Result Table 2 it clearly appears that mozzarella cheese made with an initial phospholipase treatment of the cream reduces the oiling off significantly, which is a key quality parameter for Mozzarella. The results demonstrate that it is possible
5 to obtain a stability improvement of cheese during heat treatment with the process of the invention. It is furthermore seen that the Phospholipase increases the fat content in the cheese, likely caused by a reduced fat loss during processing. Thus, the initial treatment with phospholipase leads to an increase yield in cheese production.

CLAIMS

1. A process for producing cheese, which comprises the steps of:
 - a) treating cheese milk or a fraction of the cheese
5 milk with a phospholipases; and
 - b) producing cheese from the cheese milk,
wherein step a) is conducted before and/or during step b).
2. The process of claim 1, wherein the cheese milk or the cheese milk fraction, which
10 is to be enzyme treated, comprises one or more of cream, butter or buttermilk.
3. The process of claims 1 or 2, wherein the cheese milk or the fraction of the cheese milk, which is to be enzyme-treated, comprises milk fat.
- 15 4. The process of claim 1, wherein the fraction of the cheese milk is cream.
5. The process of claim 1, wherein the fraction of the cheese milk is buttermilk.
6. The process of any of the preceding claims, wherein step a) is conducted before
20 step b).
7. The process of any of the preceding claims, wherein step a) is conducted during step b).
- 25 8. The process of claim 6, wherein the enzyme treated milk of step a) is not dried before step b).
9. The process of claim 6, wherein the enzyme treatment in step a) is conducted on a fraction of the cheese milk.
- 30 10. The process of any of the preceding claims, wherein the enzyme treatment in step a) is conducted on the cheese milk.

11. The process according to claim 9, wherein the fraction of cheese milk is cream, which process further comprises the step of subjecting the cream to pasteurization after step a) and before step b).

5

12. The process according to any of the preceding claims, further comprising the step of subjecting the cheese to a heating treatment.

13. The process according to any of the preceding claims, wherein the cheese milk
10 or the fraction of the cheese milk is subjected to a homogenization step before the production of cheese.

14. The process of claim 13, wherein the cheese is Danish Blue Cheese.

15 15. The process according to any of the preceding claims, wherein the cheese is selected from the group consisting of Campesino, Chester, Danbo, Drabant, Herregård, Manchego, Provolone, Saint Paulin, Soft cheese, Svecia, Taleggio, White cheese, Cheddar, Colby, Edam, Muenster, Gryere, Emmenthal, Camembert, Parmesan, Romano, Mozzarella, Feta; cream cheese, Neufchatel, Quarg, and
20 Queso Blanco.

16. The process according to any of the preceding claims, wherein the cheese is selected from the group consisting of rennet-curd cheese produced by rennet-coagulation of the cheese curd; ripened cheeses, such as, e.g. Cheddar, Colby,
25 Edam, Muenster, Gryere, Emmenthal, Camembert, Parmesan and Romano; fresh cheeses, such as, e.g. Mozzarella and Feta; and acid coagulated cheeses, such as, e.g., cream cheese, Neufchatel, Quarg, and Queso Blanco.

17. The process according to any of the preceding claims, further comprising the
30 step of processing the cheese into a food product, such as, e.g. pizza, ready-to-eat dishes, toast, burgers, lasagna, dressing, sauces, cheese powder, cheese flavor, processed cheese or as an ingredient in other food products.

18. The process according to any of the proceeding claims, wherein the cheese product is not processed cheese.

5 19. The process of any of the preceding claims, wherein the enzyme is selected from the group consisting of phospholipase A₁, phospholipase A₂ and phospholipase B.

20. The process of claim 19, wherein the enzyme is a phospholipase A₁ or a phospholipase A₂.

10

21. The process according to any of the preceding claims, wherein the phospholipase is an animal phospholipase or a microbial phospholipase.

22. The process of claim 21, wherein the microbial phospholipase is derived from a
15 filamentous fungi, a yeast or a bacterium.

23. The process of any of the preceding claims, wherein the phospholipase is a pancreas-derived phospholipase, such as, e.g., phospholipase A₂, e.g. from a porcine pancreas.

20

24. The process according to any of the preceding claims, wherein the phospholipase is a purified phospholipase.

25. The process according to any of the preceding claims, wherein the
25 phospholipase is provided by a *lactobacillus* used as a starter organism in the cheese production.

26. The process according to any of the proceeding claims, further comprising the step of removing or reducing the content of the phospholipase enzyme protein after
30 the enzymatic treatment.

27. The process according to any of the proceeding claims, further comprising the step of inactivating the phospholipase after the enzymatic treatment.

28. The process according to any of the preceding claims, wherein the
5 phospholipase treatment is conducted at pH 4-9, such as, e.g. pH 5-7, preferably pH 5.5-7.0.

29. The process of any preceding claim, wherein the phospholipase treatment is conducted at 3-7°C.

10

30. The process according to claim 29, wherein the enzyme treatment in step a) is conducted before step b).

31. The process according to any of claims 1-28, wherein the phospholipase
15 treatment is conducted at 30-45°C.

32. The process of claim 31, wherein the enzyme treatment is conducted during the cheese making process in step b).

20 33. The process according to any of claims 1-28, wherein the phospholipase treatment is conducted at 40-80°C, such as 45-80°C, or 50-80°C.

34. The process according to claim 33, wherein the enzyme treatment in step a) is conducted on the fraction of the cheese milk, such as, e.g. cream.

25

35. A process for producing cheese, which comprises the steps of:

a) treating a composition comprising milk fat with a phospholipase; and

b) producing cheese from the enzyme-treated composition of step a),

30 wherein step a) is conducted before step b) and/or during step b).

36. Use of a cheese produced by the process of any of the preceding claims in pizza, ready-to-eat dishes, processed cheese or as an ingredient in other food products.

- 5 37. Use of a phospholipase in the manufacturing of a cheese product, wherein the phospholipase treatment is conducted on cheese milk or a fraction thereof before and/or during the production of the cheese.

38. A cheese obtainable by the process of any of claims 1-35.

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/DK 00/00109

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A23C19/04 A23C19/032

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE FSTA 'Online! INTERNATIONAL FOOD INFORMATION SERVICE (IFIS), FRANKFURT/MAIN, DE; UMANSKII M S ET AL: "Effect on quality of Kostroma cheese of bacterial cultures selected on phospholipase activity." Database accession no. 82-1-03-p0353 XP002141695 abstract & MOLOCHNAYA PROMYSHLENNOST' 1980 VSES. NAUCHNO-ISSLED. INST. MASLODEL'NOI I SYRODEL'NOI PROMYSHLENNOSTI, UGLICH, USSR, no. 11, 1980, pages 21-25,47, ---	1-3,6-8, 10,16, 18,19, 22,28, 30-32, 35,37,38
A	US 3 156 568 A (R. HARGROVE) 10 November 1964 (1964-11-10) claim 1; examples 1,3 ---	1
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

5 July 2000

Date of mailing of the international search report

23/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Desmedt, G

INTERNATIONAL SEARCH REPORT

Int. .tional Application No

PCT/DK 00/00109

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	J. OWENS: "Observations on lecithinases from milk contaminants" PROCESS BIOCHEMISTRY, vol. 13, no. 1, 1978, page 10,12,18 XP000869553 page 12, column 2 -page 18 -----	1
A	AD. PICON: "Release of encapsulated proteinase from dehydration-rehydration liposomes by a co-encapsulated phospholipase" BIOTECHNOLOGY LETTERS, vol. 17, no. 10, 1995, pages 1051-1056, XP000870477 page 1051 -page 1053 -----	1,35
A	DATABASE WPI Section Ch, Week 198301 Derwent Publications Ltd., London, GB; Class D13, AN 1983-01331K XP002129019 & JP 57 189637 A (YAKULT HONSHA KK), 22 November 1982 (1982-11-22) abstract -----	
A	DATABASE WPI Section Ch, Week 198301 Derwent Publications Ltd., London, GB; Class D13, AN 1983-01332K XP002129020 & JP 57 189638 A (YAKULT HONSHA KK), 22 November 1982 (1982-11-22) abstract -----	
A	EP 0 531 104 A (SUMITOMO SEIKA CHEMICALS) 10 March 1993 (1993-03-10) -----	
A	US 4 612 197 A (H. POSTER) 16 September 1986 (1986-09-16) -----	
A	J. O'MAHONY: "Hydrolysis of lipoprotein fractions of milk by phospholipase C" JOURNAL OF DAIRY SCIENCE, vol. 55, no. 4, 1972, pages 408-412, XP002129018 CHAPAIN, ILLINOIS US -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. .tional Application No

PCT/DK 00/00109

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3156568 A	10-11-1964	NONE	
JP 57189637 A	22-11-1982	JP 1017655 B	31-03-1989
		JP 1533343 C	12-12-1989
JP 57189638 A	22-11-1982	JP 1017656 B	31-03-1989
		JP 1533344 C	12-12-1989
EP 531104 A	10-03-1993	CA 2077020 A	04-03-1993
		DE 69221567 D	18-09-1997
		DE 69221567 T	11-12-1997
		ES 2106146 T	01-11-1997
		JP 5268911 A	19-10-1993
		US 5478585 A	26-12-1995
US 4612197 A	16-09-1986	DE 3423699 C	16-01-1986
		AT 58459 T	15-12-1990
		CA 1233693 A	08-03-1988
		DE 3580620 D	03-01-1991
		EP 0166284 A	02-01-1986
		ES 544556 D	01-08-1986
		ES 8608296 A	01-12-1986
		PT 80711 A, B	01-07-1985
		ZA 8504431 A	26-02-1986